

=> d his

(FILE 'HOME' ENTERED AT 11:54:14 ON 10 OCT 2001)

INDEX 'ADISALERTS, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUASCI,  
BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO,  
CABA,  
CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB,  
DDFU, DGENE, DRUGB, DRUGLAUNCH, DRUGMONOG2, ...' ENTERED AT 11:54:27 ON  
10 OCT 2001

SEA (SIALIC ACID OR N-ACETYL NEURAMINIC ACID OR NANA)

-----  
80 FILE ADISALERTS  
9 FILE ADISINSIGHT  
2\* FILE ADISNEWS  
781 FILE AGRICOLA  
172 FILE ANABSTR  
442 FILE AQUASCI  
217 FILE BIOBUSINESS  
12 FILE BIOCOMMERCE  
16084 FILE BIOSIS  
362 FILE BIOTECHABS  
362 FILE BIOTECHDS  
3605 FILE BIOTECHNO  
4202 FILE CABA  
3123 FILE CANCERLIT  
17978 FILE CAPLUS  
81 FILE CEABA-VTB  
23 FILE CEN  
26 FILE CIN  
261 FILE CONFSCI  
19 FILE CROPB  
32 FILE CROPU  
584 FILE DDFB  
476 FILE DDFU  
771 FILE DGENE  
584 FILE DRUGB  
4 FILE DRUGLAUNCH  
1 FILE DRUGMONOG2  
3 FILE DRUGNL  
663 FILE DRUGU  
7 FILE DRUGUPDATES  
51 FILE EMBAL  
10977 FILE EMBASE  
2358 FILE ESBIODASE  
2 FILE FOMAD  
83 FILE FROSTI  
276 FILE FSTA  
462 FILE GENBANK  
21 FILE HEALSAFE  
362 FILE IFIPAT  
1151 FILE JICST-EPLUS  
12 FILE KOSMET  
2937 FILE LIFESCI  
2 FILE MEDICONF  
14122 FILE MEDLINE  
104 FILE NIOSHTIC  
121 FILE NTIS

133 FILE OCEAN  
 3057 FILE CASCAL  
 12 FILE CHAR  
 23 FILE PHIN  
 755 FILE PROMT  
 7993 FILE SCISEARCH  
 2 FILE SYNTHLINE  
 1399 FILE TOXLINE  
 2346 FILE TOXLIT  
 2344 FILE USPATFULL  
 835 FILE WPIDS  
 835 FILE WPINDEX

L1 QUE (SIALIC ACID OR N-ACETYL NEURAMINIC ACID OR NANA)  
 -----

FILE 'CAPLUS, BIOSIS, MEDLINE, EMBASE, SCISEARCH' ENTERED AT 11:58:39 ON  
 10 OCT 2001

L2 5867 S L1 (S) (BIOSYNTH? OR SYNTH?)  
 L3 479 S L2 (P) (ALDOLASE OR SYNTHETASE OR EPIMERASE)  
 L4 0 S L3 (P) SYNECHOCYSTIS  
 L5 44 S L3 (P) (E.COLI OR CORYNEBACTERIUM)  
 L6 19 DUP REM L5 (25 DUPLICATES REMOVED)

=> d l6 ibib ab 1-19

L6 ANSWER 1 OF 19 BIOSIS COPYRIGHT 2001 BIOSIS  
 ACCESSION NUMBER: 2001:290213 BIOSIS  
 DOCUMENT NUMBER: PREV200100290213  
 TITLE: Oligosaccharide enzyme substrates and inhibitors: methods  
 and compositions.  
 AUTHOR(S): Wong, Chi-Huey (1); Ichikawa, Yoshitaka; Shen, Gwo-Jenn  
 CORPORATE SOURCE: (1) San Diego, CA USA  
 ASSIGNEE: The Scripps Research Institute  
 PATENT INFORMATION: US 6168934 January 02, 2001  
 SOURCE: Official Gazette of the United States Patent and Trademark  
 Office Patents, (Jan. 2, 2001) Vol. 1242, No. 1, pp. No  
 Pagination. e-file.  
 ISSN: 0098-1133.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 AB Oligosaccharide compounds that are substrates and inhibitors of  
 glycosyltransferase and glycosidase enzymes and compositions containing  
 such compounds are disclosed. A method of glycosylation is also  
 disclosed.  
 An **E. coli** transformed with phagemid CMPSIL-1, which  
 phagemid comprises a gene for a modified **CMP-sialic acid**  
**synthetase** enzyme, which transformed **E. coli**  
 has the ATCC accession No. 68531 is also provided.

L6 ANSWER 2 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 1  
 ACCESSION NUMBER: 2001:527115 CAPLUS  
 TITLE: Redirection of sialic acid metabolism in genetically  
 engineered Escherichia coli  
 AUTHOR(S): Ringenber, Michael; Lichtensteiger, Carol; Vimr,  
 Eric  
 CORPORATE SOURCE: Department of Pathobiology, College of Veterinary  
 Medicine, University of Illinois at Urbana-Champaign,  
 Urbana, IL, 61802, USA  
 SOURCE: Glycobiology (2001), 11(7), 533-539  
 CODEN: GLYCE3; ISSN: 0959-6658  
 PUBLISHER: Oxford University Press  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Most microorganisms do not produce **sialic acid**  
 (sialate), and those that do appear to use a **biosynthetic**

mechanism distinct from mammals. Genetic hybrids of nonpathogenic, sialate-neg. lab. *Escherichia coli* K-12 strains designed for the de novo synthesis of the polysialic acid capsule from *E. coli* K1 proved useful in elucidating the genetics and biochem. of capsule biosynthesis. In this article we propose a dynamic model of sialometabolism to investigate the effects of biosynthetic neu (N-acetylneuraminic acid) and catabolic nan (N-acylneuraminate) mutations on the flux of intermediates through the sialate synthetic pathway. Intracellular sialate concns. were detd. by high pH anion exchange chromatog. with pulsed amperometric detection. The results indicated that a strain carrying a null defect in the gene encoding polysialyltransferase (neuS) accumulated > 50 times more CMP-sialic acid than the wild type when strains were grown in a minimal medium supplemented with glucose and casamino acids. Metabolic accumulation of CMP-sialic acid depended on a functional sialic acid synthase (neuB), as shown by the inability of a strain lacking this enzyme to accumulate a detectable endogenous sialate pool. The neuB mutant concd. trace sialate from the medium, indicating its potential value for quant. anal. of free sialic acids in complex biol. samples. The function of the sialate **aldolase** (encoded by **nanA**) in limiting intermediate flux through the **synthetic** pathway was detd. by analyzing free sialate accumulation in neuA (CMP-sialic acid synthetase) **nanA** double mutants. The combined results demonstrate how *E. coli* avoids a futile cycle in which biosynthetic sialate induces the system for its own degrdn. and indicate the feasibility of generating sialooligosaccharide precursors through targeted manipulation of sialate metab.

REFERENCE COUNT: 28  
REFERENCE(S): (1) Barrallo, S; FEBS Lett 1999, V445, P325 CAPLUS  
(2) Ferrero, M; Biochem J 1996, V317, P157 CAPLUS  
(3) Gilbert, M; Nature Biotechnol 1998, V16, P769 CAPLUS  
(4) Hoffman, B; Protein Exp Purif 1995, V6, P646 CAPLUS  
(5) Hoyer, L; Mol Microbiol 1992, V6, P873 CAPLUS  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 3 OF 19 CAPLUS COPYRIGHT 2001 ACS  
ACCESSION NUMBER: 2000:628244 CAPLUS  
DOCUMENT NUMBER: 133:218534  
TITLE: Human glycosylation enzymes and cDNAs and their use in drug screening, diagnosis, and therapy  
INVENTOR(S): Coleman, Timothy A.  
PATENT ASSIGNEE(S): Human Genome Sciences, Inc., USA  
SOURCE: PCT Int. Appl., 115 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000052136	A2	20000908	WO 2000-US5325	20000301
WO 2000052136	A3	20001228		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,  
DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 2000033884 A5 20000921 AU 2000-33884 20000301  
PRIORITY APPLN. INFO.: US 1999-122409 P 19990302  
WO 2000-US5325 W 20000301

AB The present invention relates to novel human glycosylation enzymes and isolated nucleic acids contg. the coding regions of the genes encoding such enzymes. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human glycosylation enzymes. The invention further relates to diagnostic and therapeutic methods useful for

diagnosing and treating disorders related to these novel human glycosylation enzyme polypeptides. Thus, a human cDNA encoding a protein with significant sequence homol. to mouse CMP N-acetylneuraminic acid **synthetase** was cloned and sequenced. This gene was expressed primarily in colon tissue. Another human cDNA encoded a protein with significant sequence homol. to C. jejuni cytidine **sialic acid synthetase**. A third human cDNA encoding a protein with significant sequence homol. to E. coli N-acetylneuraminic acid **aldolase** was cloned and sequenced. This gene was expressed primarily in immune cells and tissues such as primary dendritic cells, monocytes, and bone marrow.

L6 ANSWER 4 OF 19 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2000:351686 CAPLUS  
DOCUMENT NUMBER: 133:3768  
TITLE: Low cost enzymatic biosynthesis of oligosaccharides  
INVENTOR(S): Defrees, Shawn; Johnson, Karl  
PATENT ASSIGNEE(S): Neose Technologies, Inc., USA  
SOURCE: PCT Int. Appl., 103 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000029603	A2	20000525	WO 1999-US27599	19991118
WO 2000029603	A3	20001116		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 2000018261	A5	20000605	AU 2000-18261	19991118
EP 1131415	A2	20010912	EP 1999-961744	19991118

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO

PRIORITY APPLN. INFO.: US 1998-109031 P 19981118  
US 1998-109096 P 19981119  
WO 1999-US27599 W 19991118

AB This invention provides recombinant cells, reaction mixts., and methods for the enzymic synthesis of saccharides. The recombinant cells contain  
a heterologous gene that encodes a glycosyltransferase which catalyzes at least one step of the enzymic synthesis, as well a system for generating  
a nucleotide sugar that can serve as a substrate for the glycosyltransferase. The nucleotide sugar may be supplied or synthesized by an enzymic pathway comprising a sugar nucleotide regeneration cycle.

The reaction mixt. may contain a second cell type producing a nucleotide as a substrate for the sugar nucleotide regenerative cycle, preferably by a nucleotide synthase gene. Use of fusion proteins of glycosyltransferase and nucleotide sugar synthase combined with the use of an enzyme for substrate sugar synthesis is described. Chem. or enzymic sulfation may be used for the synthesis of sulfated sugars. The recombinant cells, reaction mixts., and methods are useful for efficiently synthesizing a large variety of saccharides, including polysaccharides, oligosaccharides, glycoproteins and glycolipids, using relatively low-cost starting materials. **Synthesis** of 3'-sialyllactose using **E. coli** expressing a **CMP-sialic acid synthetase**/.alpha.2,3-sialyltransferase fusion protein is described. Optional use of bakers yeast to produce CTP used in the sialic acid cycle and substrate for CMP-sialic acid synthase is also described. **Synthesis** of 3'-sialyllactose using **E. coli** expressing a **CMP-sialic acid synthetase** /.alpha.2,3-sialyltransferase fusion protein, GlcNAc 2'-**epimerase**, and **sialic acid aldolase** to **synthesize** CMP-sialic acid from GlcNAc is also described. Variations of the method using **Corynebacterium** expressing a **CMP-sialic acid synthetase** /.alpha.2,3-sialyltransferase fusion protein and CTP-**synthetase** to produce the nucleotide, nucleotide sugar, and catalyzing sugar transfer to the acceptor saccharide is described. Finally, synthesis of trisaccharide Gal.alpha.1,3Gal.beta.1,4GlcNAc using **Corynebacterium** expressing UDP-glucose pyrophosphorylase, UDP-glucose-4'-**epimerase**, .beta.1,4-galactosyltransferase, and .alpha.1,3-galactosyltransferase is described.

L6 ANSWER 5 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 2  
 ACCESSION NUMBER: 2000:202302 CAPLUS  
 DOCUMENT NUMBER: 133:115759  
 TITLE: Multiple N-acetyl neuraminic acid synthetase (neuB) genes in Campylobacter jejuni: identification and characterization of the gene involved in sialylation of lipo-oligosaccharide  
 AUTHOR(S): Linton, Dennis; Karlyshev, Andrey V.; Hitchen, Paul G.; Morris, Howard R.; Dell, Anne; Gregson, Norman A.; Wren, Brendan W.  
 CORPORATE SOURCE: Department of Neurology, United Medical and Dental School, Guy's Hospital, London, SE1 9RT, UK  
 SOURCE: Mol. Microbiol. (2000), 35(5), 1120-1134  
 CODEN: MOMIEE; ISSN: 0950-382X  
 PUBLISHER: Blackwell Science Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB N-acetylneuraminic acid (NANA) is a common constituent of Campylobacter jejuni lipo-oligosaccharide (LOS). Such structures often mimic human gangliosides and are thought to be involved in the triggering of Guillain-Barre syndrome (GBS) and Miller-Fisher syndrome (MFS) following C. jejuni infection. Anal. of the C. jejuni NCTC 11168 genome sequence identified 3 putative **NANA synthetase** genes termed neuB1, neuB2 and neuB3. The **NANA synthetase** activity of all three C. jejuni neuB gene products was confirmed by complementation expts. in an Escherichia coli neuB-deficient strain. Isogenic mutants were created in all three neuB genes, and for one such mutant (neuB1) LOS was shown to have increased mobility. C. jejuni NCTC 11168 wild-type LOS bound cholera toxin, indicating the presence of NANA in a LOS structure mimicking the ganglioside GM1. This property was lost in the neuB1

mutant. Gas chromatog.-mass spectrometry and fast atom bombardment-mass spectrometry anal. of LOS from wild-type and the neuB1 mutant strain demonstrated the lack of NANA in the latter. Expression of the neuB1

gene

in *E. coli* confirmed that NeuB1 was capable of in vitro **NANA biosynthesis** through condensation of N-acetyl-D-mannosamine and phosphoenolpyruvate. Southern anal. demonstrated that the neuB1 gene was confined to strains of *C. jejuni*

with

LOS contg. a single NANA residue. Mutagenesis of neuB2 and neuB3 did not affect LOS, but neuB3 mutants were aflagellate and non-motile. No phenotype was evident for neuB2 mutants in strain NCTC 11168, but for strain G1 the flagellin protein from the neuB2 mutant showed an apparent redn. in mol. size relative to the wild type. Thus, the neuB genes of *C. jejuni* appear to be involved in the biosynthesis of at least 2 distinct surface structures: LOS and flagella.

REFERENCE COUNT:

55

REFERENCE(S):

(1) Altschul, S; Nucleic Acids Res 1997, V25, P3389  
CAPLUS

(2) Annunziato, P; J Bacteriol 1995, V177, P312

CAPLUS

(3) Aspinall, G; Biochemistry 1994, V33, P250 CAPLUS

(4) Aspinall, G; Eur J Biochem 1993, V213, P1017  
CAPLUS

(5) Aspinall, G; Eur J Biochem 1993, V213, P1029  
CAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 3

ACCESSION NUMBER: 1999:714154 CAPLUS

DOCUMENT NUMBER: 132:1660

TITLE: Identification of Arg-12 in the active site of  
Escherichia coli K1 CMP-sialic acid synthetase

AUTHOR(S): Stoughton, Daniel M.; Zapata, Gerardo; Picone,  
Robert;

Vann, Willie F.

CORPORATE SOURCE: Laboratory of Bacterial Toxins, Division of Bacterial  
Products, OVR, CBER, FDA, Bethesda, MD, 20892, USA

SOURCE: Biochem. J. (1999), 343(2), 397-402

CODEN: BIJOAK; ISSN: 0264-6021

PUBLISHER: Portland Press Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Escherichia coli K1 CMP-sialate **synthetase** (I) catalyzes the **synthesis** of CMP-sialic acid from CTP and **sialic acid**. The active site of 418-amino-acid *E. coli* I was localized to its N-terminal half. The bacterial I enzymes have a conserved motif, IAIIPARXXSKGLXXKN, at their N-termini. Several basic residues were identified at or near the active site of the *E. coli* enzyme by chem. modification and site-directed mutagenesis. Only 1 of the Lys residues in the N-terminal motif, Lys-21, appeared to be essential for activity. Mutation of Lys-21 in the N-terminal motif resulted in an inactive enzyme. Furthermore, Arg-12 of the N-terminal motif appeared to be an active site residue, based on the following evidence. Substituting Arg-12 with Gly or Ala resulted in inactive enzymes, indicating that this residue is required

for

enzymic activity. The R12K mutant was partially active, demonstrating that a pos. charge is required at this site. Steady-state kinetic anal. revealed changes in kcat, Km, and Ks values for CTP, which implicated Arg-12 in catalysis and substrate binding.

REFERENCE COUNT:

25

REFERENCE(S):

(1) Ambrose, M; Biochemistry 1992, V31, P775 CAPLUS

(2) Caligiuri, M; J Biol Chem 1991, V266, P8328

CAPLUS

(4) Ganguli, S; J Bacteriol 1994, V176, P4583 CAPLUS

- L6 ANSWER 7 OF 19 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 1998:415986 BIOSIS  
DOCUMENT NUMBER: PREV199800415986  
TITLE: Characterization of the CTP binding sites in the **E. coli K1 CMP-sialic acid synthetase.**  
AUTHOR(S): Stoughton, D. M.; Vann, W. F.  
CORPORATE SOURCE: Food Drug Administration, Bethesda, MD USA  
SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (1998) Vol. 98, pp. 125.  
Meeting Info.: 98th General Meeting of the American Society for Microbiology Atlanta, Georgia, USA May 17-21, 1998  
American Society for Microbiology  
. ISSN: 1060-2011.  
DOCUMENT TYPE: Conference  
LANGUAGE: English
- L6 ANSWER 8 OF 19 SCISEARCH COPYRIGHT 2001 ISI (R)  
ACCESSION NUMBER: 1998:408767 SCISEARCH  
THE GENUINE ARTICLE: ZK302  
TITLE: Arginine-12 is essential for CTP binding and catalysis of the **E. coli K1 CMP-sialic acid synthetase**  
AUTHOR: Stoughton D M (Reprint); Zapata G; Vann W F  
CORPORATE SOURCE: FDA, CBER, BETHESDA, MD  
COUNTRY OF AUTHOR: USA  
SOURCE: FASEB JOURNAL, (31 JUL 1997) Vol. 11, No. 9, Supp. [S], pp. 3303-3303.  
Publisher: FEDERATION AMER SOC EXP BIOL, 9650 ROCKVILLE PIKE, BETHESDA, MD 20814-3998.  
ISSN: 0892-6638.  
DOCUMENT TYPE: Conference; Journal  
FILE SEGMENT: LIFE  
LANGUAGE: English  
REFERENCE COUNT: 0
- L6 ANSWER 9 OF 19 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 1997:422568 BIOSIS  
DOCUMENT NUMBER: PREV199799721771  
TITLE: Arginine-12 is essential for CTP binding and catalysis of the **E. coli K1 CMP-sialic acid synthetase.**  
AUTHOR(S): Stoughton, D. M.; Zapata, G.; Vann, W. F.  
CORPORATE SOURCE: CBER, FDA, Bethesda, MD USA  
SOURCE: FASEB Journal, (1997) Vol. 11, No. 9, pp. A1421.  
Meeting Info.: 17th International Congress of Biochemistry and Molecular Biology in conjunction with the Annual Meeting of the American Society for Biochemistry and Molecular Biology San Francisco, California, USA August 24-29, 1997  
ISSN: 0892-6638.  
DOCUMENT TYPE: Conference; Abstract  
LANGUAGE: English
- L6 ANSWER 10 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 4  
ACCESSION NUMBER: 1997:508112 CAPLUS  
DOCUMENT NUMBER: 127:216834  
TITLE: Purification and characterization of the Escherichia coli K1 neuB gene product N-acetylneuraminic acid synthetase  
AUTHOR(S): Vann, Willie F.; Tavarez, Jose J.; Crowley, Jane;

CORPORATE SOURCE: Vimr, Eric; Silver, Richard P.  
Laboratory of Bacterial Polysaccharides, Center for  
Biologics Research and Review, Bethesda, MD, 20892,  
USA  
SOURCE: Glycobiology (1997), 7(5), 697-701  
CODEN: GLYCE3; ISSN: 0959-6658  
PUBLISHER: Oxford University Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Escherichia coli K1 produces a capsular polysaccharide of  
.alpha.(2-8)-poly-N-acetylneuraminic acid. This polysaccharide is an  
essential virulence factor of these neuropathogenic bacteria. The genes  
necessary for the synthesis of neuNAc were localized to a plasmid contg.  
the neuBAC genes of the K1 gene cluster. Cells harboring the neuB+  
allele  
in an **aldolase** (nanA-) neg. background produce neuNAc in vivo.  
Enzymic synthesis of neuNAc could be demonstrated in exts. of cells  
harboring an expression plasmid (pNEUB) contg. the neuB gene alone.  
NeuNAc **synthetase** was purified to homogeneity from exts. of  
cells harboring pNEUB. The mol. wt. of the purified enzyme is 40 kDa,  
similar to that predicted by the nucleotide sequence of the neuB gene.  
The N-terminal sequence of the purified protein matches that predicted by  
the nucleotide sequence of the neuB gene. NeuNAc **synthetase**  
catalyzes the formation of neuNAc as indicated by its coupling to the  
CMP-NeuNAc **synthetase** reaction. The enzyme condenses ManNAc and  
PEP with the release of phosphate. The **E.coli** neuNAc  
**synthetase** is specific for ManNAc and PEP, unlike rat liver enzyme  
that utilizes N-acetylmannosamine-6-phosphate to form NeuNAc-9-PO4. This  
represents the first report of a purifn. of a **sialic**  
**acid synthetase** from either a eukaryotic or prokaryotic  
source to homogeneity. These expts. clearly demonstrate an  
**aldolase-independent sialic acid**  
**synthetase** activity in **E. coli** K1.

L6 ANSWER 11 OF 19 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1997:486007 CAPLUS  
TITLE: Chemo-enzymic synthesis and structural analysis of  
polysialic acid.  
AUTHOR(S): McGowen, M. M.; Jennings, H. J.; Vann, W. F.  
CORPORATE SOURCE: CBER, FDA, Bethesda, MD, 20892, USA  
SOURCE: Book of Abstracts, 214th ACS National Meeting, Las  
Vegas, NV, September 7-11 (1997), CARB-051. American  
Chemical Society: Washington, D. C.  
CODEN: 64RNAO  
DOCUMENT TYPE: Conference; Meeting Abstract  
LANGUAGE: English

AB Poly-.alpha.(2,8)-N-acetyl neuraminic acid (polysialic acid, PSA) has  
been  
implicated in the pathogenic mechanism of Escherichia coli K1 urinary  
tract infection and neonatal meningitis. The three dimensional structure  
of PSA is rationalized as random coil with localized helixes of n = 9-11.  
The carboxylate group is essential to stabilizing the active conformation  
of PSA. In this study we are investigating the role that the hydroxyl  
moiety at C9 plays. Our approach was to **synthesize**  
**sialic acid** analogs modified at C9 by chem. methods and  
incorporate analogs into polymer by using enzymes. 9-Azidosialic acid  
was  
charged with CMP using recombinant CMP-NeuNAc **synthetase**.  
Mutant **E. coli** K1, EV241, which can neither  
**synthesize** nor degrade **sialic acid** was the  
source of sialyltransferase. Incubation of the activated azido-monomer  
with total membrane preps. of EV241 enzymically synthesized modified  
.alpha.(2,8)-PSA de novo. The 9-modified PSA was isolated and purified  
by  
gel filtration and anion exchange chromatog. The homopolysaccharide was  
then confirmed by <sup>13</sup>C NMR spectroscopy and Dionex integrated amperometry



HPLC anal. The data presented may assist in the further understanding of the active conformation of PSA involved in pathogenesis or immunological response.

L6 ANSWER 12 OF 19 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 5  
ACCESSION NUMBER: 1996:154855 BIOSIS  
DOCUMENT NUMBER: PREV199698726990  
TITLE: Characterization of cpsF and its product  
CMP-N-acetylneuraminic acid synthetase, a group B streptococcal enzyme that can function in K1 capsular polysaccharide biosynthesis in Escherichia coli.  
AUTHOR(S): Haft, Rachel F.; Wessels, Michael R. (1); Mebane, Mary Fisk; Conaty, Neil; Rubens, Craig E.  
CORPORATE SOURCE: (1) Channing Lab., Brigham Women's Hosp., Div. Infectious Diseases, Beth Israel Hosp., Harvard Med. Sch., 180 Longwood Avenue, Boston, MA 02115 USA  
SOURCE: Molecular Microbiology, (1996) Vol. 19, No. 3, pp. 555-563.

ISSN: 0950-382X.

DOCUMENT TYPE: Article  
LANGUAGE: English

AB Group B Streptococcus (GBS) is the foremost cause of neonatal sepsis and meningitis in the United States. A major virulence factor for GBS is its capsular polysaccharide, a high molecular weight polymer of branched oligosaccharide subunits. N-acetylneuraminic acid (Neu5Ac or **sialic acid**), at the end of the polysaccharide side chains, is critical to the virulence function of the capsular polysaccharide. Neu5Ac must be activated by CMP-Neu5Ac **synthetase** before it is incorporated into the polymer. We showed previously that a transposon mutant of a serotype III GBS strain which had no detectable capsular Neu5Ac was deficient in CMP-Neu5Ac-**synthetase** activity (Wessels et al., 1992). In this paper, we report the identification and characterization of cpsF, a gene interrupted by transposon insertion in the previously described Neu5Ac-deficient mutant. The predicted amino acid

sequence of the cpsF gene product shares 57% similarity and 37% identity with CMP-Neu5Ac **synthetase** encoded by the Escherichia coli K1 gene, neuA. The enzymatic function of the protein encoded by cpsF was established by cloning the gene in *E. coli* under the control of the T7 polymerase/promoter. Lysates of *E. coli* in which the cpsF gene product was expressed, catalysed the condensation of CTP with Neu5Ac to form CMPNeu5Ac. In addition, when a CMP-Neu5Ac **synthetase**-deficient mutant of *E. coli* K1 was transformed with cpsF, K1 antigen expression was restored. We conclude that cpsF encodes CMP-Neu5Ac **synthetase** in type III GBS, and that the GBS enzyme can function in the capsule-**synthesis** of a heterologous bacterial species.

L6 ANSWER 13 OF 19 MEDLINE DUPLICATE 6  
ACCESSION NUMBER: 96150083 MEDLINE  
DOCUMENT NUMBER: 96150083 PubMed ID: 8579837  
TITLE: CMP-N-acetyl neuraminic-acid synthetase from Escherichia coli: fermentative production and application for the preparative synthesis of CMP-neuraminic acid.  
AUTHOR: Kittelmann M; Klein T; Kragl U; Wandrey C; Ghisalba O  
CORPORATE SOURCE: Ciba-Geigy Ltd, Pharmaceuticals Division, Basel, Switzerland.  
SOURCE: APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, (1995 Dec) 44 (1-2) 59-67.  
PUB. COUNTRY: GERMANY: Germany, Federal Republic of  
LANGUAGE: English  
FILE SEGMENT: B  
ENTRY MONTH: 199603

(X)

ENTRY DATE: Entered STN: 19960327  
Last Updated on STN: 19980206  
Entered Medline: 19960321

AB In an optimized sorbitol/yeast extract/mineral salt medium up to 12 U/l  
**CMP-N-acetyl-neuraminic-acid**  
(Neu5Ac) **synthetase** was produced by Escherichia coli K-235 in  
shake-flask culture. A colony mutant of this strain, **E.**  
**coli** K-235/CS1, was isolated with improved enzyme formation: in  
shake flasks with a yield of up to 20.8 U/l and 54 mU/mg protein in the  
cell extract. With this strain 26500 U CMP-Neu5Ac **synthetase** was  
produced with a high specific activity (0.128 U/mg) by fed-batch  
fermentation on 230-l scale. On a 10-l scale the enzyme yield was 191 U/l  
culture medium. The enzyme was partially purified by precipitation with  
polyethyleneglycol resulting in a three- to fourfold enrichment and a  
recovery rate of more than 80%; most of the CTP hydrolysing enzymes were  
removed. The native **synthetase** was deactivated completely by  
incubation at 45 degrees C for 10 min, but could be stabilized remarkably  
by glycerol and different salts. The enzyme was used for the preparative  
**synthesis** of CMP-Neu5Ac with a conversion yield of 87% based on  
CTP.

L6 ANSWER 14 OF 19 CAPLUS COPYRIGHT 2001 ACS  
ACCESSION NUMBER: 1993:146291 CAPLUS  
DOCUMENT NUMBER: 118:146291  
TITLE: CMP-sialic acids manufacture with microbial cell  
extracts  
INVENTOR(S): Kittelmann, Matthias; Ghisalba, Oreste; Klein,  
Teresa;  
PATENT ASSIGNEE(S): Kragl, Udo; Wandrey, Christian Prof Dr  
Ciba-Geigy A.-G., Switz.; Forschungszentrum Juelich  
GmbH  
SOURCE: Eur. Pat. Appl., 25 pp.  
CODEN: EPXXDW  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 524143	A1	19930120	EP 1992-810522	19920708
EP 524143	B1	19971210		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, PT, SE				
AT 161051	E	19971215	AT 1992-810522	19920708
ES 2110481	T3	19980216	ES 1992-810522	19920708
CA 2073954	AA	19930118	CA 1992-2073954	19920715
AU 9220348	A1	19930121	AU 1992-20348	19920716
AU 664036	B2	19951102		
JP 05276973	A2	19931026	JP 1992-189647	19920716
IL 102527	A1	19960804	IL 1992-102527	19920716
US 5334514	A	19940802	US 1993-152269	19931112
PRIORITY APPLN. INFO.:			CH 1991-2119	A 19910717
			US 1992-915474	B1 19920716

AB **CMP-sialic acids** are prepd. by incubation of CTP and  
**sialic acids** with microbial cell exts. contg.  
cytidine-5'-monophospho-N-acetylneuraminic acid **synthetase**  
activity. Escherichia coli was cultured and an ext. was prepd. which was  
used to prep. CMP-Neu5Ac from CTP and N-acetylneuraminic acid (Neu5Ac).  
Methods for optimizing **E. coli** growth and enzyme yield  
and for further purifn. of the enzyme were described. An **E.**  
**coli** mutant with higher yields of the enzyme was produced.

L6 ANSWER 15 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 7  
ACCESSION NUMBER: 1992:470187 CAPLUS  
DOCUMENT NUMBER: 117:70187  
TITLE: Overproduction of CMP-sialic acid synthetase for

organic synthesis  
AUTHOR(S): Liu, Jennifer Lin Chun; Shen, Jenn; Ichikawa, Yoshitaka; Rutan, James F.; Zapata, Gerardo; Vann, Willie F.; Wong, Chi Huey  
CORPORATE SOURCE: Dep. Chem., Scripps Res. Inst., La Jolla, CA, 92037, USA  
SOURCE: J. Am. Chem. Soc. (1992), 114(10), 3901-10  
CODEN: JACSAT; ISSN: 0002-7863  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The gene coding for Escherichia coli CMP-**sialic acid synthetase** (E.C. 2.7.7.43) was cloned and overexpressed in **E. coli** through a primer-directed polymerase chain reaction. Two plasmids were constructed to produce the native enzyme and a modified enzyme fused with a decapeptide at the C-terminus. The decapeptide tag was used for detection of the enzyme prodn. Both enzymes produced from **E. coli** were isolated and purified with an orange A dye resin and FPLC. Contrary to the native enzyme, the modified enzyme is more active at higher pH. Studies on specificity indicate that both enzymes have a high specific activity for C-9 modified NeuAc derivs. at neutral pH. Some C-5 modified (hydroxy-, deoxy-, and deoxyfluoro-) NeuAc derivs. are not acceptable as substrates. The modified enzyme has been used in the synthesis of CMP-NeuAc from ManNAc and CMP and sialyl N-acetyllactosamine (Neu.alpha.2,6Gal.beta.1,4GlcNAc) with in situ generation of NeuAc and regeneration of CMP-NeuAc. The 6-O-acyl derivs. of ManNAc were prepd. via transesterification in anhyd. DMF by using an engineered stable subtilisin variant as a catalyst, and the products were used as substrates in **sialic acid aldolase**-catalyzed **synthesis** of 9-O-acyl-NeuAc derivs.

L6 ANSWER 16 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 8  
ACCESSION NUMBER: 1986:17473 CAPLUS  
DOCUMENT NUMBER: 104:17473  
TITLE: Regulation of sialic acid metabolism in Escherichia coli: role of N-acylneuraminate pyruvate-lyase  
AUTHOR(S): Vimr, Eric R.; Troy, Frederic A.  
CORPORATE SOURCE: Sch. Med., Univ. California, Davis, CA, 95616, USA  
SOURCE: J. Bacteriol. (1985), 164(2), 854-60  
CODEN: JOBAAY; ISSN: 0021-9193  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB In **E. coli**, **synthesis** of **sialic acid** is not regulated by allosteric inhibition mediated by cytidine 5'-monophospho-N-acetylneuraminic acid (CMP-NeuNAc). Evidence for the lack of metabolic control by feedback inhibition was demonstrated by measuring the intracellular levels of **sialic acid** and CMP-NeuNAc in mutants defective in **sialic acid** polymn. and in CMP-NeuNAc **synthesis**. Polymn.-defective mutants could not **synthesize** the poly(**sialic acid**) capsule and accumulated 25-fold more CMP-NeuNAc than the wild type. Mutants unable to activate **sialic acid** because of a defect in CMP-NeuNAc **synthetase** accumulated .apprx.7-fold more **sialic acid** than the wild type. An addnl. 3-fold increase in **sialic acid** levels occurred when a mutation resulting in loss of N-acylneuraminate pyruvate-lyase (**sialic acid aldolase**) was introduced into the CMP-NeuNAc **synthetase**-deficient mutant. The **aldolase** mutation could not be introduced into the polymn.-defective mutant, suggesting that any further increase in the intracellular CMP-NeuNAc concn. was toxic. Thus, sialic acid **aldolase** can regulate the intracellular concn. of sialic acid and, therefore, the concn. of CMP-NeuNAc. Thus, regulation of **aldolase**, mediated by sialic acid induction, is necessary not only for dissimilating sialic acid but also for modulating the level of metabolic intermediates in the sialic acid pathway. In agreement with this conclusion, an increase in the intracellular sialic acid concn. was correlated with an increase in

**aldolase** activity. Direct evidence for the central role of **aldolase** in regulating the metabolic flux of sialic acid in *E. coli* was that exogenous, radiolabeled sialic acid was specifically incorporated into sialyl polymer in an **aldolase**-neg. strain, but not in the wild type.

L6 ANSWER 17 OF 19 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 9  
ACCESSION NUMBER: 1986:3295 CAPLUS  
DOCUMENT NUMBER: 104:3295  
TITLE: Identification of an inducible catabolic system for sialic acids (nan) in *Escherichia coli*  
AUTHOR(S): Vimr, Eric R.; Troy, Frederic A.  
CORPORATE SOURCE: Sch. Med., Univ. California, Davis, CA, 95616, USA  
SOURCE: J. Bacteriol. (1985), 164(2), 845-53  
CODEN: JOBAAY; ISSN: 0021-9193  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB *Escherichia coli* K-12 and K-12 hybrid strains constructed to express a polysialic acid capsule, the K1 antigen, were able to efficiently use sialic acid as a sole C source. This ability was dependent on induction of *gtoreq.2* activities: a sialic acid-specific transport activity, and an **aldolase** activity specific for cleaving sialic acids. Induction over basal levels required sialic acid as the apparent inducer, and induction of both activities was repressed by glucose. Induction also required the intracellular accumulation of **sialic acid**, which could be either added exogenously to the medium or accumulated intracellularly through **biosynthesis**. Exogenous sialic acid appeared to be transported by an active mechanism that did not involve covalent modification of the sugar. Mutations affecting either the transport or degradn. of sialic acid prevented its use as a C source and have been designated *nanT* and *nanA*, resp. These mutations were located by transduction near *min 69* on the *E. coli* K-12 genetic map, between *argG* and *glnF*. In addn. to being unable to use sialic acid as a C source, **aldolase**-neg. mutants were growth-inhibited by this sugar. Therefore, the intracellularly accumulated sialic acid was toxic in **aldolase**-deficient *E. coli* strains. The dual role of **aldolase** in dissimilating and detoxifying sialic acids is consistent with the apparent multiple controls on expression of this enzyme.

L6 ANSWER 18 OF 19 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
ACCESSION NUMBER: 84202093 EMBASE  
DOCUMENT NUMBER: 1984202093  
TITLE: Purification and properties of *E. coli* cytidine 5'-monophosphate **n-acetyl-neuraminic acid synthetase**.  
AUTHOR: Vann W.F.; Kotsatos M.; Chang K.; et al.  
CORPORATE SOURCE: National Center for Drugs and Biologics, FDA, Bethesda, MD 20205, United States  
SOURCE: Federation Proceedings, (1984) 43/6 (no. 1644).  
CODEN: FEPA7  
COUNTRY: United States  
DOCUMENT TYPE: Journal  
FILE SEGMENT: 029 Clinical Biochemistry  
LANGUAGE: English

L6 ANSWER 19 OF 19 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 1985:245274 BIOSIS  
DOCUMENT NUMBER: BA79:25270  
TITLE: PURIFICATION AND PROPERTIES OF N ACETYLNEURAMINATE LYASE EC-4.1.3.3 FROM *ESCHERICHIA-COLI*.  
AUTHOR(S): UCHIDA Y; TSUKADA Y; SUGIMORI T  
CORPORATE SOURCE: KYOTO RES. LAB., MARUKIN SHOYU CO., LTD., UJI, KYOTO 611.  
SOURCE: J BIOCHEM (TOKYO), (1984) 96 (2), 507-522.

FILE SEGMENT: BA 001D  
LANGUAGE: English

AB N-Acetylneuraminic lyase [N-acetylneuraminic acid **aldolase** EC 4.1.3.3] from *E. coli* was purified by protamine sulfate treatment, fractionation with  $(\text{NH}_4)_2\text{SO}_4$ , column chromatography

on

DEAE-Sephacel, gel filtration on Ultrogel AcA 44 and preparative polyacrylamide gel electrophoresis. The purified enzyme preparation was homogenous on analytical polyacrylamide gel electrophoresis, and was free from contaminating enzymes including NADH oxidase and NADH dehydrogenase. The enzyme catalyzed the cleavage of N-acetylneuraminic acid to N-acetylmannosamine and pyruvate in a reversible reaction. Both cleavage and **synthesis** of N-acetylneuraminic acid had the same pH optimum .apprx. 7.7. The enzyme was stable at pH 6.0-9.0, and was thermostable up to 60.degree. C. The thermal stability increased up to 75.degree. C in

the

presence of pyruvate. No metal ion was required for the enzyme activity, but heavy metal ions such as  $\text{Ag}^+$  and  $\text{Hg}^{2+}$  were potent inhibitors.

Oxidizing agents such as N-bromosuccinimide, I, and  $\text{H}_2\text{O}_2$ , and

SH-inhibitors

such as P-chloromercuribenzoic acid and  $\text{HgCl}_2$  were also potent

inhibitors.

The  $K_m$  values for N-acetylneuraminic acid and N-glycolylneuraminic acid were 3.6 mM and 4.3 mM, respectively. Pyruvate inhibited the cleavage reaction competitively;  $K_i$  was calculated to be 1.0 mM. In the condensation reaction, N-acetylglucosamine, N-acetylgalactosamine, glucosamine and galactosamine could not replace N-acetylmannosamine as substrate, and phosphoenolpyruvate, lactate, .beta.-hydroxypyruvate and other pyruvate derivatives could not replace pyruvate as substrate. The

MW

of the native enzyme was estimated to be 98,000 by gel filtration

methods.

After denaturation in sodium dodecyl sulfate or in 6 M guanidine-HCl, the MW was reduced to 33,000, indicating the existence of 3 identical subunits. The enzyme could be used for the enzymatic determination of **sialic acid**; reaction conditions were devised for determining the bound form of **sialic acid** by coupling neuraminidase from *Arthrobacter ureafaciens*, lactate dehydrogenase and NADH.

=> d his

(FILE 'HOME' ENTERED AT 14:42:25 ON 10 OCT 2001)

INDEX 'ADISALERTS, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUASCI,  
BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO,  
CABA,  
CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB,  
DDFU, DGENE, DRUGB, DRUGLAUNCH, DRUGMONOG2, ...' ENTERED AT 14:42:38 ON  
10 OCT 2001

SEA (N-ACETYLGLUCOSAMINE 2'-EPIMERASE) OR (UDP-GLCNAC

2'-EPIMER

-----  
0\* FILE ADISNEWS  
5 FILE BIOSIS  
1 FILE CANCERLIT  
3 FILE CAPLUS  
1 FILE CONFSCI  
4 FILE EMBASE  
1 FILE LIFESCI  
4 FILE MEDLINE  
3 FILE SCISEARCH  
1 FILE TOXLIT

L1 QUE (N-ACETYLGLUCOSAMINE 2'-EPIMERASE) OR (UDP-GLCNAC

2'-EPIMER

-----  
FILE 'BIOSIS, EMBASE, MEDLINE, CAPLUS, SCISEARCH' ENTERED AT 14:47:25 ON  
10 OCT 2001

L2 19 S L1

L3 6 DUP REM L2 (13 DUPLICATES REMOVED)

=> d l3 ibib ab 1-6

L3 ANSWER 1 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 1  
ACCESSION NUMBER: 1999:311054 BIOSIS  
DOCUMENT NUMBER: PREV199900311054  
TITLE: Mutations in the human UDP-N-acetylglucosamine 2-epimerase  
gene define the disease sialuria and the allosteric site  
of

the enzyme.  
AUTHOR(S): Seppala, Raili; Lehto, Veli-Pekka; Gahl, William A. (1)  
CORPORATE SOURCE: (1) NICHD, NIH, 10 Center Drive, Building 10, Room 9S-241,  
Bethesda, MD, 20892-1830 USA  
SOURCE: American Journal of Human Genetics, (June, 1999) Vol. 64,  
No. 6, pp. 1563-1569.  
ISSN: 0002-9297.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AB Sialuria is a rare inborn error of metabolism characterized by  
cytoplasmic  
accumulation and increased urinary excretion of free N-acetylneuraminic  
acid (NeuAc, sialic acid). Overproduction of NeuAc is believed to result  
from loss of feedback inhibition of uridinediphosphate-N-  
**acetylglucosamine 2-epimerase (UDP-  
GlcNAc 2-epimerase)** by cytidine  
monophosphate-N-acetylneuraminic acid (CMP-Neu5Ac). We report the cloning  
and characterization of human UDP-GlcNAc 2-epimerase cDNA, with mutation  
analysis of three patients with sialuria. Their heterozygote mutations,  
R266W, R266Q, and R263L, indicate that the allosteric site of the  
epimerase resides in the region of codons 263-266. The heterozygous  
nature  
of the mutant allele in all three patients reveals a dominant mechanism  
of  
inheritance for sialuria.

L3 ANSWER 2 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 2  
ACCESSION NUMBER: 1999:397525 BIOSIS  
DOCUMENT NUMBER: PREV199900397525  
TITLE: UDP-GlcNAc 2-epimerase: A regulator of cell surface  
sialylation.  
AUTHOR(S): Keppler, Oliver T.; Hinderlich, Stephan; Langner, Josmar;  
Schwartz-Albiez, Reinhard; Reutter, Werner; Pawlita,  
Michael (1)  
CORPORATE SOURCE: (1) Applied Tumor Virology Program, Deutsches  
Krebsforschungszentrum, Im Neuenheimer Feld 280, D-69120,  
Heidelberg Germany  
SOURCE: Science (Washington D C), (May 21, 1999) Vol. 284, No.  
5418, pp. 1372-1376.  
ISSN: 0036-8075.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AB Modification of cell surface molecules with sialic acid is crucial for  
their function in many biological processes, including cell adhesion and  
signal transduction. Uridine diphosphate-N-  
**acetylglucosamine 2-epimerase (UDP-  
GlcNAc 2-epimerase)** is an enzyme that  
catalyzes an early, rate-limiting step in the sialic acid biosynthetic  
pathway. UDP-GlcNAc 2-epimerase was found to be a major determinant of

cell surface sialylation in human hematopoietic cell lines and a critical regulator of the function of specific cell surface adhesion molecules.

L3 ANSWER 3 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 3  
ACCESSION NUMBER: 1997:486001 BIOSIS  
DOCUMENT NUMBER: PREV199799785204  
TITLE: A bifunctional enzyme catalyzes the first two steps in N-acetylneuraminic acid biosynthesis of rat liver: Molecular cloning and functional expression of UDP-N-acetyl-glucosamine 2-epimerase/N-acetylmannosamine kinase.  
AUTHOR(S): Staesche, Roger; Hinderlich, Stephan; Weise, Christoph; Effertz, Karin; Lucka, Lothar; Moormann, Petra; Reutter, Werner (1)  
CORPORATE SOURCE: (1) Inst. Molekularbiol. Biochemie, Freie Univ. Berlin, Arnimallee 22, D-14195 Berlin-Dahlem Germany  
SOURCE: Journal of Biological Chemistry, (1997) Vol. 272, No. 39, pp. 24319-24324.  
ISSN: 0021-9258.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
AB N-Acetylneuraminic acid (Neu5Ac) is the precursor of sialic acids, a group of important molecules in biological recognition systems. Biosynthesis of Neu5Ac is initiated and regulated by its key enzyme, UDP-N-acetylglucosamine 2-epimerase (UDP-GlcNAc 2-epimerase, EC 5.1.3.14)/N-acetylmannosamine kinase (ManNAc kinase, EC 2.7.1.60) in rat liver (Hinderlich, S., Stasche, R., Zeitler, R., and Reutter, W. (1997) J. Biol. Chem. 272, 24313-24318). In the present paper we report the isolation and characterization of a cDNA clone encoding this bifunctional enzyme. An open reading frame of 2166 base pairs encodes 722 amino acids with a predicted molecular mass of 79 kDa. The deduced amino acid sequence contains exact matches of the sequences of five peptides derived from tryptic cleavage of the enzyme. The recombinant bifunctional enzyme was expressed in COS7 cells, where it displayed both epimerase and kinase activity. Distribution of UDP-GlcNAc 2-epimerase/ManNAc kinase in the cytosol of several rat tissues was investigated by determining both specific enzyme activities. Secreting organs (liver, salivary glands, and intestinal mucosa) showed high specific activities of UDP-GlcNAc 2-epimerase/ManNAc kinase, whereas significant levels of these activities were absent from other organs (lung, kidney, spleen, brain, heart, skeletal muscle, and testis). Northern blot analysis revealed no UDP-GlcNAc 2-epimerase/ManNAc kinase mRNA in the non-secreting tissues.

L3 ANSWER 4 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 4  
ACCESSION NUMBER: 1997:486000 BIOSIS  
DOCUMENT NUMBER: PREV199799785203  
TITLE: A bifunctional enzyme catalyzes the first two steps in N-acetylneuraminic acid biosynthesis of rat liver: Purification and characterization of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase.  
AUTHOR(S): Hinderlich, Stephan; Staesche, Roger; Zeitler, Reinhard; Reutter, Werner (1)  
CORPORATE SOURCE: (1) Inst. Molekularbiol. Biochemie, Freie Univ. Berlin, Arnimallee 22, D-14195 Berlin-Dahlem Germany  
SOURCE: Journal of Biological Chemistry, (1997) Vol. 272, No. 39, pp. 24313-24318.  
ISSN: 0021-9258.  
DOCUMENT TYPE: Article  
LANGUAGE: English  
AB Biosynthesis of N-acetylneuraminic acid (Neu5Ac), a prominent component of glycoconjugates, is initiated by the action of UDP-N-acetylglucosamine 2-epimerase (UDP-



**GlcNAc 2-epimerase**, EC 5.1.3.14) and N-acetylmannosaminyl kinase (ManNAc kinase, EC 2.7.1.40). We demonstrate for the first time that the two activities are part of one bifunctional enzyme in rat liver. The enzyme was purified to homogeneity from rat liver cytosol using salmine sulfate precipitation and chromatography on phenyl-Sepharose, ATP-agarose, and Mono Q. The purification resulted in one polypeptide with an apparent molecular mass of 75 kDa. Immunoprecipitation with a polyclonal antibody against the polypeptide reduced both enzyme activities in equal amounts. Gel filtration analysis of purified UDP-GlcNAc 2-epimerase/ManNAc kinase showed that the polypeptide self-associates as a dimer and as a hexamer with apparent molecular masses of 150 and 450 kDa, respectively. The hexamer was fully active for both enzyme activities, whereas the dimer catalyzed only the phosphorylation of N-acetylmannosamine (ManNAc). Incubation of the dimer with UDP-N-acetylglucosamine led to reassembly of the fully active hexamer; maximal quantities of the hexamer were produced after incubation for 3 h. Kinetic analysis of purified hexameric and dimeric enzyme revealed significantly lower Michaelis constants ( $93 \pm 3$  to  $121 \pm 15$   $\mu$ M for ManNAc and  $1.18 \pm 0.13$  to  $1.67 \pm 0.20$  mM for ATP) and higher cooperativity (Hill coefficients of  $1.42 \pm 0.16$  to  $1.17 \pm 0.06$  for ManNAc and  $1.30 \pm 0.09$  to  $1.05 \pm 0.14$  for ATP) for the hexamer for both substrates of ManNAc kinase. The Michaelis constant of UDP-GlcNAc 2-epimerase for its substrate was  $11 \pm 2$   $\mu$ M. The Hill coefficient of  $0.45 \pm 0.07$  represents strongly negative cooperativity in substrate binding. UDP-GlcNAc 2-epimerase was feedback inhibited by CMP-Neu5Ac. Complete inhibition was achieved with 60  $\mu$ M CMP-Neu5Ac, and highly positive cooperativity (Hill coefficient of 4.1) was found for inhibitor binding.

L3 ANSWER 5 OF 6 SCISEARCH COPYRIGHT 2001 ISI (R)

ACCESSION NUMBER: 1998:76055 SCISEARCH

THE GENUINE ARTICLE: YQ995

TITLE: Molecular cloning of the rate-limiting enzyme in sialic acid synthesis, uridinediphosphate-N-acetylglucosamine-2-epimerase (UDP-GlcNAc-2-epimerase): Implications for sialuria.

AUTHOR: Seppala R (Reprint); Lehto V P; Gahl W A

CORPORATE SOURCE: NICHD, SECT HUMAN BIOCHEM GENET, HERITABLE DISORDERS BRANCH, NIH, BETHESDA, MD; UNIV OULU, DEPT PATHOL, OULU, FINLAND

COUNTRY OF AUTHOR: USA; FINLAND

SOURCE: AMERICAN JOURNAL OF HUMAN GENETICS, (OCT 1997) Vol. 61, No. 4, Suppl. [S], pp. 1519-1519.  
 Publisher: UNIV CHICAGO PRESS, 5720 S WOODLAWN AVE, CHICAGO, IL 60637.  
 ISSN: 0002-9297.

DOCUMENT TYPE: Conference; Journal

FILE SEGMENT: LIFE; CLIN

LANGUAGE: English

REFERENCE COUNT: 0

L3 ANSWER 6 OF 6 BIOSIS COPYRIGHT 2001 BIOSIS

ACCESSION NUMBER: 1998:111440 BIOSIS

DOCUMENT NUMBER: PREV199800111440

TITLE: Molecular cloning of the rate-limiting enzyme in sialic acid synthesis, uridinediphosphate-N-acetylglucosamine-2-epimerase (UDP-GlcNAc-2-epimerase): Implications for sialuria.

AUTHOR(S): Seppala, R. (1); Lehto, V. P.; Gahl, W. A.

CORPORATE SOURCE: (1) Section Human Biochem. Genetics, Heritable Disorders Branch, NICHD, National Inst. Health, Bethesda, MD USA

SOURCE: American Journal of Human Genetics, (Oct., 1997) Vol. 61, No. 4 SUPPL., pp. A261.

Meeting Info.: 47th Annual Meeting of the American Society  
of Human Genetics Baltimore, Maryland USA October  
28- November 1, 1997  
ISSN: 0002-9297.

DOCUMENT TYPE:  
LANGUAGE:

Conference  
English